

QuickVue[®]

Product Catalog QuickVue Rapid Lateral-Flow Tests





Proven testing for faster diagnosis and treatment

Healthcare professionals have relied on our QuickVue products for more than 30 years. Test accurately and quickly for a range of infectious diseases, women's health conditions, and many other diseases and conditions.

QuidelOrtho's industry-leading expertise in immunoassays, molecular testing, clinical chemistry and transfusion medicine delivers fast, accurate and reliable diagnostics to diverse healthcare settings. Partnering globally, we pioneer innovations, forging seamless connections and illuminating paths to informed decisions.



QuickVue menu of assays

QuickVue Influenza A+B Test

CLIA waived

Immunoassay for the differential detection of influenza type A and type B from nasal swab and nasopharyngeal swab (NPS) specimens, direct and viral transport medium (VTM).

- Results in 10 minutes or less
- Three steps NPS
- Less than 30 seconds hands-on time
- One reagent
- High specificity for influenza
- Not affected by variations of seasonal influenza strains

Catalog #	Quantity	Description
20183	25 tests	QuickVue Influenza A+B test kit
20103	25 swabs	Nasal swab pack
0171	25 swabs	Influenza nasal swab transport tubes
20226	100 swabs	Nasopharyngeal swab pack

QuickVue SARS Antigen Test

Immunoassay for the qualitative detection of SARS-CoV-2 antigen in anterior nares specimens.

- Results in 10 minutes
- 96.6% positive percent agreement
- 99.3% negative percent agreement
- One reagent
- No age restriction
- Available with Emergency Use Authorization only

Catalog #	Quantity	Description
20387	25 tests	QuickVue SARS Antigen test kit

QuickVue RSV Test

CLIA waived

Immunoassay for the qualitative detection of respiratory syncytial virus (RSV) from nasopharyngeal swab, nasopharyngeal aspirate or nasal/nasopharyngeal wash specimens for symptomatic pediatric patients (18 years of age and younger).

- Results in 15 minutes or less
- 99% sensitivity with nasopharyngeal aspirate
- 92% sensitivity with nasopharyngeal swab
- 83% sensitivity with nasal/nasopharyngeal wash
- Three-step nasopharyngeal swab, one-step nasopharyngeal aspirate and nasal/nasopharyngeal wash with approximately 30 seconds hands-on time
- One reagent

Catalog #	Quantity	Description
20193	20 tests	QuickVue RSV test kit

QuickVue Adenoviral Conjunctivitis Test

CLIA waived

Immunoassay for the visual, qualitative in vitro detection of adenoviral antigens (hexon protein) directly from human eye fluid.

- Results in 10 minutes
- 90% sensitivity and 96% specificity
- Simple four-step procedure
- Disposable test with no additional equipment needed to administer or interpret
- Reduce inappropriate antibiotic use

Catalog #	Quantity	Description
20324	10 tests	QuickVue adenoviral conjunctivitis test kit
RPS-ADSTD	1	QuickVue adenoviral conjunctivitis external controls

QuickVue+® Strep A Test

Immunoassay for the rapid qualitative detection of group A streptococcal antigen from throat swab specimens.

- Results in approximately 5 minutes
- Plus/minus (+/-) readout
- 98% overall agreement with sheep blood agar culture
- Two-reagent procedure to reduce hands-on time
- Test complete indicator lets you read the test at your convenience
- CLIA Moderately Complex

Catalog #	Quantity	Description
20122	25 tests	QuickVue+ Strep A test kit
00354	1	Strep A liquid control set
20227	50 swabs	Strep A swabs

QuickVue In-Line® Strep A Test

CLIA waived

Unique in-line extraction to detect group A streptococcal antigen directly from patient throat swab specimens.

- Results in 5 minutes or less
- 96% overall agreement with SBA culture
- One easy step via in-line extraction
- Single reagent method allows for less chance of operator error
- Individually packaged for convenient single-use testing and disposal
- First CLIA-waived strep A test

Catalog #	Quantity	Description
00343	25 tests	QuickVue in-line strep A test kit
00345	6 Pos/6 Neg	QuickVue in-line strep A control set
00347	75 swabs	Strep A swab pack

QuickVue Dipstick Strep A Test

CLIA waived

Immunoassay for the rapid qualitative detection of group A streptococcal antigen from throat swab specimens.

- Results in 5 minutes or less
- 95% overall agreement with SBA culture
- High sensitivity at low levels of colony count

Catalog #	Quantity	Description
20108	50 tests	QuickVue dipstick strep A test kit
00354	1	Strep A liquid control set
20227	50 swabs	Strep A swabs

QuickVue+ hCG Combo Test

Immunoassay for the qualitative detection of human chorionic gonadotropin (hCG) in serum or urine for the early detection of pregnancy.

- Results in 5 minutes (serum), 3 minutes (urine)
- Plus/minus (+/-) readout
- >99% accuracy
- 10 mIU/mL sensitivity, serum; 20 mIU/mL sensitivity, urine
- Detects low hCG levels as early as 2-3 days before expected menses
- CLIA waived when using urine
- CLIA Moderately Complex when using serum


Catalog #	Quantity	Description
00178	30 tests	QuickVue+ hCG combo test kit
00179	90 tests	QuickVue+ hCG combo test kit
00272	1	hCG control set – urine
00281	1	hCG control set – serum

QuickVue hCG Combo Test

Immunoassay for the qualitative detection of hCG in serum or urine for the early detection of pregnancy.

- Results in 5 minutes (serum), 3 minutes (urine)
- >99% accuracy
- 25 mIU/mL sensitivity in serum and urine
- Detects low hCG levels as early as 2-3 days before expected menses
- CLIA waived when using urine
- CLIA Moderately Complex when using serum

Catalog #	Quantity	Description
20110	50 tests	QuickVue hCG combo test Kit
00272	1	hCG control set – urine
00281	1	hCG control set – serum



QuickVue hCG Urine Test

CLIA waived

Immunoassay for the qualitative detection of hCG in urine for the early detection of pregnancy.

- Results in 3 minutes or less
- >99% accuracy
- 25 mIU/mL sensitivity
- Detects low hCG levels as early as 2-3 days before expected menses

Catalog #	Quantity	Description
20109	25 tests	QuickVue hCG urine test kit
00272	1	hCG control set - urine

QuickVue TLI H. pylori Stool Antigen Test

Rapid membrane enzyme immunoassay for the qualitative detection of *Helicobacter pylori* specific antigen in a single-use cassette.

- Results in 30 minutes
- 97% sensitivity; 100% specificity
- Demonstrate loss of antigen following treatment
- Can be used with various sample types including fresh, frozen, Cary Blair or culture and sensitivity (C&S) samples
- CLIA Moderately Complex

Catalog #	Quantity	Description
2043	25 tests	QuickVue TLI H. pylori stool antigen test kit

QuickVue TLI Campylobacter Test

Rapid membrane enzyme-linked immunosorbent assay for the qualitative detection of a *Campylobacter*-specific antigen in human fecal specimens.

- Results in 30 minutes
- >97% sensitivity; >99% specificity
- Does not rely on bacterial viability
- Can be performed with fresh, frozen, Cary Blair or C&S samples or specimens
- CLIA Moderately Complex

Catalog #	Quantity	Description
20344	25 tests	QuickVue TLI Campylobacter test kit

QuickVue TLI Lactoferrin Test

Immunochemical test for the qualitative detection of elevated levels of lactoferrin, a marker for fecal leukocytes and an indicator of intestinal inflammation.

- Results in 10 minutes
- >100% sensitivity; >97% specificity
- Can be used with samples stored at room temperature for up to 2 weeks
- CLIA Moderately Complex

Catalog #	Quantity	Description
20345	25 tests	QuickVue TLI lactoferrin test kit

QuickVue H. pylori Test

Immunoassay for the rapid qualitative detection of immunoglobulin G (IgG) antibodies specific to *Helicobacter pylori*.

- Results in 5 minutes or less
- 92% overall agreement with enzyme immunoassay
- Simple, one-step procedure from only two hanging drops of whole blood, or one drop of serum or plasma
- CLIA waived when using whole blood specimens
- CLIA Moderately Complex when using serum or plasma

Catalog #	Quantity	Description
0W009	10 tests	QuickVue H. pylori test kit
0W010	30 tests	QuickVue H. pylori test kit

QuickVue iFOB Test

CLIA waived

(*A Fecal Immunochemical or FIT test*) Immunochemical device intended for the qualitative detection of fecal occult blood.

- Results in 5-10 minutes
- Analytically more sensitive than guaiac-based tests*
- Analytical sensitivity: 50 ng hHb/mL buffer or 50 µg hHb/g feces
- Specific to human hemoglobin
- No dietary restrictions
- Only one specimen collection required

Catalog #	Quantity	Description
20194	20 tests	QuickVue iFOB test kit
20196	10 tests	Patient specimen collection kit
20201	50 tests	QuickVue iFOB tray pack (test cassettes)
20204	40 tests	Patient specimen collection kit
20205	100 tests	Test cassettes
20197	1	iFOB control set

*Park DI, Ryu S, Kim YH, et al. Comparison of guaiac-based and quantitative immunochemical fecal occult blood testing in a population at average risk undergoing colorectal cancer screening. *Am J Gastroenterol.* 2010;105(9):2017-2025. doi:10.1038/ajg.2010.179



9975 Summers Ridge Road
San Diego, CA 92121 USA
800-874-1517
quidelortho.com

New QuidelOrtho branding may not be available in all markets, subject to country-specific regulatory approval. Please confirm with your local commercial team.

©2024 QuidelOrtho Corporation.
All trademarks referenced are trademarks of their respective owners.

PR-105926-CRP-EN-US-v1